## **Post-Surgical Instructions for Open Wounds**

- 1. Please avoid showering and leave all bandages/dressings intact and in place for the next 2 days.
- 2. Only if the dressing falls off or becomes severely saturated, you may remove the dressings and begin wound care as follows:

Cleanse the area with warm tap water using a cotton swab or gauze pad

You may shower normally at this time

Dry the area with a cotton swab or gauze pad then apply Polysporin/Bacitracin or petrolatum along with a new clean bandaid or non-stick dressing and paper tape.

DO NOT USE NEOSPORIN OR TRIPLE ANTIBIOTIC

Change the dressing daily as above UNTIL THE WOUND IS COMPLETELY HEALED

- 1. You may alternate Extra Strength Tylenol with Ibuprofen (Advil) as needed for discomfort. If alternating the medications, they should be taken at least 4 hrs. apart.
- 2. Continue taking all medications prescribed by your physician including aspirin if you are already taking it.
- 3. Apply an ice pack over the surgical site for 20 minutes at a time, two to three times a day of surgery to decrease pain, swelling, and bruising that may occur.
- 4. Elevate the area if possible to reduce swelling over the next 48 hours after surgery on two pillows. It is normal to have swelling and bruising around the surgical site.
- 5. No strenuous activity for 48 hours.
- 6. Continue all other prescribed medication.
- 7. Do not drink alcoholic beverages for 48 hours. Do not smoke for 3 weeks as smoking is detrimental to wound healing.
- 8. Avoid activities that require you to bend at the waist and elevate your blood pressure as this can increase pressure and bleeding.

## **BLEEDING:**

- 1. Leave the bandage in place and use rolled up gauze or a clean cloth to apply firm, direct pressure over the surgical site for 20 minutes.
- 2. Reapply pressure for an additional 20 minutes if necessary followed by 20 minutes of ice.
- 3. Call the on-call physician at the number below if pressure and ice fails to stop the bleeding
- 4. Use additional gauze and tape to reinforce bandage once the bleeding has stopped.

## In Case of an EMERGENCY please call our office immediately at (315) 452-3376, Press option #1 to reach the on-call provider.

Antibiotic:	Take one by mouth once every 12 hours for
10 days	

REPAIR:

Date:

Initial: